

IN THE CLAIMS:

1. (Currently amended) A computer-implemented method for calculating a global hydrophobic moment of a tertiary protein structure comprising a plurality of residues, wherein the method is run on a system comprising one or more distinct devices, each of
5 the one or more distinct devices being embodied on a tangible computer-readable recordable storage medium, the method comprising:

calculating a centroid of residue centroids, wherein calculating the centroid of residue centroids is carried out by a tertiary protein structure analyzer executing on a computer configured to carry out the step of calculating the centroid;

10 using the centroid of residue centroids as a spatial origin of a global linear hydrophobic moment, wherein using the centroid of residue centroids as a spatial origin of a global linear hydrophobic moment is carried out by a tertiary protein structure analyzer executing on a computer configured to carry out the step of using the centroid as a spatial origin;

15 calculating a first-order hydrophobic moment, wherein calculating the first-order hydrophobic moment is carried out by a tertiary protein structure analyzer executing on a computer configured to carry out the step of calculating the first-order hydrophobic moment;

enhancing correlation between residue centroid magnitude and residue
20 solvent accessibility, wherein the correlation between residue centroid magnitude and residue solvent accessibility is enhanced using a distance metric, wherein enhancing correlation between residue centroid magnitude and residue solvent accessibility is carried out by a tertiary protein structure analyzer executing on a computer configured to carry out the step of enhancing correlation;

25 using the first-order hydrophobic moment and the enhanced correlation between residue centroid magnitude and residue solvent accessibility to ~~define~~ calculate the global linear hydrophobic moment, wherein each of the residue centroids contributes a magnitude and direction to the global linear hydrophobic moment, and wherein using the first-order hydrophobic moment and the enhanced correlation between residue
30 centroid magnitude and residue solvent accessibility to calculate the global linear hydrophobic moment comprises measuring a fractional distance of each residue centroid

to [a] an ellipsoidal surface of the tertiary protein structure ~~is measured~~, and mapping
each ellipsoidal coordinate onto a sphere with radius equal to a major principal axis if a
residue centroid has a same fractional distance from a center of the protein structure to
the ellipsoidal surface of the tertiary protein structure as one or more additional residue
5 centroids, wherein mapping places each residue at a same distance from the center of the
protein structure to enable that each such residue that has the same fractional distance
from the center of the protein structure contributes an equivalent magnitude to the global
linear hydrophobic moment wherein each residue centroid having a same fractional
distance to a surface of the tertiary protein structure as one or more additional residue
10 centroids contributes an equivalent magnitude to the global linear hydrophobic moment
as the one or more additional residue centroids by mapping each residue at a same
distance from a center of the protein structure, wherein using the first-order hydrophobic
moment and the enhanced correlation between residue centroid magnitude and residue
solvent accessibility to ~~define~~ calculate the global linear hydrophobic moment is carried
15 out by a tertiary protein structure analyzer executing on a computer configured to carry
out the step of using the first-order hydrophobic moment and the enhanced correlation
between residue centroid magnitude and residue solvent accessibility to ~~define~~ calculate
the global linear hydrophobic moment;

using the global linear hydrophobic moment to characterize an
20 amphiphilicity of a tertiary protein structure, wherein using the global linear hydrophobic
moment to characterize the amphiphilicity of the tertiary protein structure is carried out
by a tertiary protein structure analyzer executing on a computer configured to carry out
the step of using the global linear hydrophobic moment to characterize the amphiphilicity
of the tertiary protein structure; and

25 outputting the global linear hydrophobic moment to a user.

2. (Canceled)

3. (Original) The method of claim 1, wherein the correlation between residue
30 centroid magnitude and residue solvent accessibility is enhanced using an ellipsoidal
metric.

4. (Original) The method of claim 1, wherein the correlation between residue centroid magnitude and residue solvent accessibility is enhanced using a solvent accessibility metric.

5

5. (Original) The method of claim 1, wherein the centroid of residue centroids represents a geometric center of the tertiary protein structure.

6. (Cancelled)

10

7. (Original) The method of claim 1, wherein the global linear hydrophobic moment characterizes a magnitude of amphiphilicity of the tertiary protein structure.

8. (Original) The method of claim 1, wherein the global linear hydrophobic moment characterizes a direction of amphiphilicity of the tertiary protein structure.

15

9. (Original) The method of claim 1, wherein the global linear hydrophobic moment is used to identify functional regions of the tertiary protein structure.

20 10. (Cancelled)

11. (Cancelled)

12. (Cancelled)

25

13. (Cancelled)

14. (Currently amended) An apparatus for calculating a global hydrophobic moment of a tertiary protein structure comprising a plurality of residues, the apparatus comprising:

30

a memory; and

at least one processor operative to:

calculate a centroid of residue centroids;

use the centroid of residue centroids as a spatial origin of a global linear hydrophobic moment;

5 calculate a first-order hydrophobic moment;

enhance correlation between residue centroid magnitude and residue solvent accessibility, wherein the correlation between residue centroid magnitude and residue solvent accessibility is enhanced using a distance metric;

use the first-order hydrophobic moment and the enhanced correlation
10 between residue centroid magnitude and residue solvent accessibility to ~~define~~ calculate
the global linear hydrophobic moment, wherein each of the residue centroids contributes
a magnitude and direction to the global linear hydrophobic moment, and wherein using
the first-order hydrophobic moment and the enhanced correlation between residue
15 centroid magnitude and residue solvent accessibility to calculate the global linear
hydrophobic moment comprises measuring a fractional distance of each residue centroid
to [a] an ellipsoidal surface of the tertiary protein structure ~~is measured~~, and mapping
each ellipsoidal coordinate onto a sphere with radius equal to a major principal axis if a
residue centroid has a same fractional distance from a center of the protein structure to
the ellipsoidal surface of the tertiary protein structure as one or more additional residue
20 centroids, wherein mapping places each residue at a same distance from the center of the
protein structure to enable that each such residue that has the same fractional distance
from the center of the protein structure contributes an equivalent magnitude to the global
linear hydrophobic moment ~~wherein each residue centroid having a same fractional~~
~~distance to a surface of the tertiary protein structure as one or more additional residue~~
25 ~~centroids contributes an equivalent magnitude to the global linear hydrophobic moment~~
~~as the one or more additional residue centroids by mapping each residue at a same~~
~~distance from a center of the protein structure;~~

use the global linear hydrophobic moment to characterize an amphiphilicity of a tertiary protein structure; and

30 output the global linear hydrophobic moment to a user.

15. (Original) The apparatus of claim 14, wherein the centroid of the residue centroids represents a geometric center of the tertiary protein structure.

16. (Cancelled)

5

17. (Original) The apparatus of claim 14, wherein the global linear hydrophobic moment is used to identify functional regions of the tertiary protein structure.

18. (Canceled)

10

19. (Original) The apparatus of claim 14, wherein the correlation between residue centroid magnitude and residue solvent accessibility is enhanced using an ellipsoidal metric.

15 20. (Original) The apparatus of claim 14, wherein the correlation between residue centroid magnitude and residue solvent accessibility is enhanced using a solvent accessibility metric.

21. (Currently amended) An article of manufacture for calculating a global hydrophobic moment of a tertiary protein structure comprising a plurality of residues, comprising:

20 a computer-readable medium having computer-readable code embodied thereon, the computer-readable code comprising:

25 a step for calculating a centroid of residue centroids;
a step for using the centroid of residue centroids as a spatial origin of a global linear hydrophobic moment;

a step for calculating a first-order hydrophobic moment;
a step for enhancing correlation between residue centroid magnitude and residue solvent accessibility, wherein the correlation between residue centroid magnitude and residue solvent accessibility is enhanced using a distance metric;

30 a step for using the first-order hydrophobic moment and the enhanced

correlation between residue centroid magnitude and residue solvent accessibility to ~~define~~
calculate the global linear hydrophobic moment, wherein each of the residue centroids
contributes a magnitude and direction to the global linear hydrophobic moment, and
wherein using the first-order hydrophobic moment and the enhanced correlation between
5 residue centroid magnitude and residue solvent accessibility to calculate the global linear
hydrophobic moment comprises measuring a fractional distance of each residue centroid
to [a] an ellipsoidal surface of the tertiary protein structure is measured, and mapping
each ellipsoidal coordinate onto a sphere with radius equal to a major principal axis if a
residue centroid has a same fractional distance from a center of the protein structure to
10 the ellipsoidal surface of the tertiary protein structure as one or more additional residue
centroids, wherein mapping places each residue at a same distance from the center of the
protein structure to enable that each such residue that has the same fractional distance
from the center of the protein structure contributes an equivalent magnitude to the global
linear hydrophobic moment ~~wherein each residue centroid having a same fractional~~
15 ~~distance to a surface of the tertiary protein structure as one or more additional residue~~
~~centroids contributes an equivalent magnitude to the global linear hydrophobic moment~~
~~as the one or more additional residue centroids by mapping each residue at a same~~
~~distance from a center of the protein structure;~~
a step for using the global linear hydrophobic moment to characterize an
20 amphiphilicity of a tertiary protein structure; and
a step for outputting the global linear hydrophobic moment to a user.